



FEVER



What is fever?

- ❖ Fever is your body's normal reaction to an infection – it is a good thing
- ❖ Fever is a temperature of greater than 100.4 degrees Fahrenheit in children under 2 months
- ❖ Fever is a temperature greater than 101 degrees in children older than 2 months of age.
- ❖ A temperature less than 100.4 is normal and does not need to be treated.

What is the best way to take a temperature?

- ❖ Rectal temperatures are the most accurate and should always be used in babies under 2 months
- ❖ An underarm temperature in children greater than 2 months of age may be used. You do not need to add or subtract a degree.
- ❖ Ear thermometers are notoriously inaccurate.

Should I be afraid of a fever? NO!!

- ❖ Fever will not cause brain damage and will not hurt your child whether it is 101 or 104. Remember, fever is a good thing. It is your body fighting an infection.
- ❖ How your child looks, and feels is more important than the number on the thermometer. Bring the fever down, then assess how your child is feeling.

Why should I treat a fever?

- ❖ While your child has a fever, they may feel achy and lethargic.
- ❖ Treating the fever may make your child feel better in the interim, although it will not make the infection go away.

What medicines can I use to treat a fever?

- ❖ Tylenol is the brand name for acetaminophen
- ❖ Motrin and Advil are brand names for ibuprofen Children less than 6 months of age can be given acetaminophen (Tylenol) every 4 hours as needed.
- ❖ Children over 6 months of age can be giving Tylenol every 4 hours or Ibuprofen (Motrin/Advil) every 6 hours as needed.



Westchester Commons

15400 WC Commons Way
Midlothian, VA 23113-7320
Tel: (804) 549-5405
Fax: (804) 379-8162

Old Jahnke Road

7023 Old Jahnke Road
Richmond, VA 23225-4126
Tel: (804) 320-1353
Fax: (804) 320-6636

Harbour Pointe

6510 Harbour View Court, Suite 100
Midlothian, VA 23112-6559
Tel: (804) 739-8166
Fax: (804) 639-6614



COMMONWEALTH PEDIATRICS

Parent With Confidence

- ❖ Tylenol and Motrin/Advil are two different medicines. They shouldn't be given at the same time. Tylenol needs to be given 4 hours from Tylenol.
- ❖ Motrin/Advil needs to be given 6 hours from Motrin/Advil. Because they are different medicines, they can be given closer together. For example, you can give Tylenol then 2-3 hours later give Motrin/Advil then 2-3 hours later give Tylenol...
- ❖ These medicines should only be given when your child actually has a fever.
- ❖ Tylenol and Ibuprofen take approximately 30 minutes to work. Give the medicine time to work.
- ❖ Please take a Tylenol and ibuprofen dosage form at the front desk the next time you visit the office.

Call our office if:

- ❖ any child less than 2 months of age has a fever
- ❖ fever persists longer than 72 hours
- ❖ fever does not respond to Tylenol or ibuprofen
- ❖ fever greater than 105

Pediatric Advisor 2006.4; Copyright © 2006 McKesson Corporation and/or one of its subsidiaries. All Rights Reserved. Written by B.D. Schmitt, M.D., author of "Your Child's Health," Bantam Books. This content is reviewed periodically and is subject to change as new health information becomes available. The information is intended to inform and educate and is not a replacement for medical evaluation, advice, diagnosis or treatment by a healthcare professional.



Westchester Commons

15400 WC Commons Way
Midlothian, VA 23113-7320
Tel: (804) 549-5405
Fax: (804) 379-8162

Old Jahnke Road

7023 Old Jahnke Road
Richmond, VA 23225-4126
Tel: (804) 320-1353
Fax: (804) 320-6636

Harbour Pointe

6510 Harbour View Court, Suite 100
Midlothian, VA 23112-6559
Tel: (804) 739-8166
Fax: (804) 639-6614